

Methodology Article

Investigation on Major Cause of Organ Condemnation and Its Economic Significance in Cattle Slaughtered in Chiro Municipal Abattoir, Ethiopia

Bayan Ahmed

Department of Veterinary Medicine, Haramaya University, Haramaya, Ethiopia

Email address:

drbayan2017@gmail.com

To cite this article:

Bayan Ahmed. Investigation on Major Cause of Organ Condemnation and Its Economic Significance in Cattle Slaughtered in Chiro Municipal Abattoir, Ethiopia. *Animal and Veterinary Sciences*. Vol. 11, No. 4, 2023, pp. 87-93. doi: 10.11648/j.avs.20231104.11

Received: February 7, 2023; **Accepted:** July 3, 2023; **Published:** July 26, 2023

Abstract: A cross sectional study was carried out from November 2021 to April 2022 with aim to identify the major cause of organ condemnation, associated risk factors and to estimate direct financial losses attributed to the condemned organ in cattle slaughtered at Chiro Municipal abattoir, Eastern Ethiopia. Prior to slaughter, animals were subjected to ante and post mortem examination to identify physical abnormalities and gross pathological lesions responsible for organ condemnation. During ante mortem examination, in 75 (19.3) cattle various types of abnormalities were detected include; lacrimation (2.6%), pale mucus membrane (3.09%), nasal discharge (2.9%), hernia (1%), salivation (0.8%), lameness (1.8%), emaciation (0.41%), depression (1%), blindness (0.41%), local swelling (0.27%) and rough hair coat (2%). Out of 384 cattle examined 185 (48.2%) were animals positive from which different organ condemned during postmortem inspection. Among organs examined, 105 (27.3%) livers, 55 (14.3%) lungs, 18 (4.68%) hearts and 7 (1.8%) kidneys were condemned. Major pathological conditions that caused a total condemnation of organs were fasciolosis (14.6%), hydatidosis (9.38%), cirrhosis (6.25%), calcification (2.6%), emphysema (2.34%), pericarditis (2.6%), hydronephrosis (1.7%), and pneumonia (4.09%) *Cysticercus bovis* (0.5%) and abscess (1.56%). Organ condemnation rates show significant difference ($P < 0.05$) among age group, origin of Animals and status of body condition. The total financial loss estimated in this study, due to organ condemnation was 138,784.00 ETB per annum. Therefore, the observation of such level of abnormalities and substantial financial loss with condemnation of affected organs warrants the veterinary institution to design appropriate control measures.

Keywords: Abattoir, Cattle, Chiro, Ethiopia, Financial Loss, Organs Condemnation

1. Introduction

Ethiopia is a developing nation with the most livestock in Africa, with an estimated population of 53.99 million cows, 25.5 million sheep, 24.06 million goats, 1.91 million horses, 6.75 million donkeys, 0.35 million mules, 0.92 million camels, and 50.38 million chickens. [12] Ethiopia's cattle industry formerly made a sizable contribution to the national economy, and it continues to hold out hope for the future of the nation's economy. About 52.13 million cattle heads are present in the nation.

In the entirety of Sub-Saharan Africa, animal production has traditionally been considered the route to agricultural development. Domesticated animals are fundamental to

Ethiopia's economy, like they are in many other developing countries. They provide food in the form of meat and milk as well as non-food products including transportation services, draft power, manure, and other inputs for the cultivation of food crops and fuel for cooking. The selling of the aforementioned goods as well as animal hides and skins is another way that livestock generate money. Ethiopia, which is the first in Africa and tenth in the world, is well recognized for having a large cattle population. According to latest animal population estimates, the nation has roughly 1.9 million horses, 6.75 million sheep, 49.56 million goats, and 53.99 million heads of cattle. [15]

Ethiopia has a significant quantity of cattle, however the industry is known for its low output. Low genetic potential of

native breeds, poor management, poor diet, poor reproductive ability, high illness incidence, and parasite burden are all contributing factors to the low output. In addition, each year a sizeable loss is caused by animal deaths, inadequate weight growth, and the rejection of edible parts and carcasses at the time of slaughter during regular meat inspection. The cattle business suffers a productivity loss that is expected to cost more than \$900 million a year [25].

The monitoring of numerous human and animal illnesses was greatly aided by abattoirs. All animals entering the human food chain can be checked at the slaughterhouse for strange symptoms, lesions, or particular illnesses. Diseases caused by parasites, bacteria, and viruses are the main reason for organ condemnation during post-mortem examination. [4]. One method of determining the illness state of a herd has been identified as monitoring and other conditions at slaughter; however, this source of data is not widely utilized globally. The incidence and epidemiology of animal illness problems may be determined using data from abattoirs, which can also be used to calculate the financial losses resulting from the condemnation of afflicted organs [29].

In order to identify animal products with aberrant pathological lesions that are unsightly and unfit for human consumption, meat inspection is done in the slaughterhouse. Meat inspection helps identify specific livestock illnesses, stops the spread of contaminated meat that might cause disease in humans and animals, and ensures product competitiveness in the local market [21]. The infections caused by parasites, bacteria, and viruses are the major reasons for organ condemnation during post-mortem examination. The primary issues are liver flukes and hydatid cysts in the lung, liver, and kidney [31].

Tropical parasites cost the meat industry more than any other ailment by a wide margin by kidanu, [26]. Ethiopia has conducted several researches to identify the main illness conditions seen during ante mortem and post mortem inspection and to estimate the economic significance of organ

condemnation, just like many other tropical nations [24]. Therefore, in order to build an effective plan for preventive and controls, it would be important to have thorough information on the occurrence of various causes and their economic loss. The research area currently lacks slaughterhouse surveillance data on the prevalence of different diseases and economic loss from organ condemnation. Therefore, the objective of present study was:

- 1) To determine the main factors that lead to organ condemnation in cattle slaughtered at the Chiro Municipal Abattoir.
- 2) To determine the potential risk factors in the study area that could be related to organ condemnation.
- 3) To calculate the financial damage directly attributable to the organs of condemned animals that was slaughtered at Chiro Municipal Abattoir.

2. Materials and Methods

2.1. Study Area Description

The study was conducted in Chiro municipal abattoir from November, 2021 to April; 2022. Chiro is a capital town of West Hararge zone, which is located 326km far away from capital city of Ethiopia, Finfinnee. The district has 40 rural peasant associations and one urban administration. The weather climate of the woredas is characterized by four main seasons in a year. The dry April and long rainy season that extend from July to the end of November. The district has daily mean a temperature ranging from 27.5-38.5°C and rain fall ranging from 900-1800mm. The agro ecological zone of the district can be categorized into, high land which portion 8%, midland 38% and lowland 54%. Its altitude ranges between 2500m and 3748m above sea level. The livestock populations of the district are cattle 92656, goat 67962, sheep 37594, horse 403, mule 234, donkey 15404, camel 626 and poultry 105,253. [11].




Figure 1. Map of study area.

2.2. Study Population

The research animals were cattle from several woreda in and around Chiro, such Oda Bultum, Gemachis, Tulo, Mieso, and others, which were transported to the town abattoir for slaughter. These animals were kept in an extensive production system, and both cars and people were used to convey them to the slaughterhouse. Animals were chosen using a simple random sample procedure, regardless of their body condition scores, sex, or age.

2.3. Study Design

From November 2020 to April 2021, a cross-sectional study was conducted in order to identify the primary reason for organ condemnation and to estimate the direct economic loss incurred as a result of organ condemnation in cattle slaughtered at the Chiro Municipal abattoir.

2.4. Sample Size and Sampling Method

Since no previous research had been done in the area, the following parameters were used to calculate the necessary sample size: 95% level of confidence (CL), 5% desirable level of accuracy, and 50% prevalence. The formula provided by Thrusfield, [42], was used to calculate the sample size.

$$n = \frac{1.96^2 \times \text{Pexp} (1 - \text{Pexp})}{d^2}$$

Where:

n= the total sample size;

Pexp= expected prevalence;

d= absolute precision

Accordingly, the calculated sample size was 384. Thus 384 Cattle were considered for ante and post mortem examination.

2.5. Study Methodology

2.5.1. Ante-Mortem Inspection

Cattle were examined at the lairage before to slaughter to determine their age, sex, physical condition score, provenance, and any anomalies they may have had. Based on tooth eruption trends, two traditional age groups—young 2 to 6 years and adult >6 years—were created by Pace and Wakeman [38]. According to the approach described by Nicholson and Butterworth [34], the body condition of the cattle was scored and divided into three categories: poor, medium, and good.

2.5.2. Post Mortem Inspection

Each visceral organ, particularly the liver, lung, heart, and kidney, was thoroughly examined postmortem by vision, inspection, palpation, and systematic incision for the presence of cysts, different adult parasites, and other abnormalities. The guidelines for meat inspection were used to differentiate and evaluate the pathological lesions [16].

2.6. Financial Assessment

Based on the average number of animals slaughtered in the

slaughterhouse annually from retrospective data of the abattoir, the condemnation rate of each kind of studied organ, and the condemnation rate of each organ, the total financial loss due to organ condemnation was calculated. From the butcheries in Chiro town, the average local market price of each organ was obtained through a questionnaire. The following formula developed by Ogunrinade and Ogunrinade [37], in 1980, was used to determine the total direct financial loss:

$$EL = \Sigma srx \times Coy \times Roz$$

Where:

EL = Annual economic loss estimated due to organ condemnation.

Σsrx = Annual number of cattle slaughtered at the abattoir.

Coy = Average cost of each liver/lung/heart/kidney.

Roz = Condemnation rate of each liver/ lung/heart/kidney.

2.7. Data Analysis

For data analysis, SPSS software version 16.0 was utilized. Collected data were put on a specifically designed form, coded, and entered. The level of condemnation, expressed as the ratio of condemned organs to all investigated organs, was calculated using descriptive statistics. Pearson's Chi-square was used to analyze the differences between condemnation rates, age, sex, and bodily condition ratings (X2). When the p-value for this study's outcome is less than 0.05, it was deemed statistically significant.

3. Results

Out of 384 cattle examined upon ante mortem examination, various types of abnormalities were detected on 75 (19.3) of cattle. The abnormalities detected include: localized lesion, hernia, lacrimation, nasal discharge, rough hair coat, lameness, depression, blindness (Table 1).

Table 1. Rate of abnormalities encountered during ante mortem inspection in cattle slaughtered at Chiro municipal abattoir.

Abnormality	No of affected animals	Percentage (%)
Localized lesion	21	5.5
Hernia	4	1
Lacrimation	10	2.6
Nasal discharge	11	2.9
Rough hair coat	8	2
Lameness	7	1.8
Depression	4	1
Salivation	3	0.8
Total	75	19.3

In the Chiro municipal abattoir, 384 cattle were examined during the study period. Of the inspected organs, 185 (48.2%) were positive for various lesions detected on various organs, among which 72 (18.8%) and 113 (29.4%) were condemned for various reasons. Young cattle 61 (32.9%) were more likely to be condemned than mature cattle 124 (67.1%), and male cattle 159 (85.9%) were more likely to be condemned than female cattle 26 (14.1%). Medium body condition cattle had

the highest condemnation rates, followed by good and poor cattle with scores of 101 (54.6%), 56 (30.3%), and 28 (15.5%), respectively. Cattle from OdaBultum share the largest percentage, 80 (43.2%), of the 185 (48.2%) condemned rates,

followed by Tulo 49 (26.5%), Gemachis 27 (14.6%), and Mieso worda 29 (15.7%), in that order. Age, bodily condition ratings, and origin all varied statistically significantly ($P < 0.05$) when possible risk variables were examined in animal models.

Table 2. Prevalence of affected organ with regard to Sex, Age, BCS and origins.

Variables	No. of examined	Cattle	Positive (%)	X ²	P-value
Sex	Male	333	159 (85.9%)	0.478	0.490
	Female	51	26 (14.1%)		
Age	Young	151	61 (32.9%)	6.441	0.011
	Adult	233	124 (67.1%)		
BCS	Poor	32	28 (15.5%)	25.203	0.000
	Medium	207	101 (54.6%)		
Origin	Good	145	56 (30.3%)	10.506	0.014
	Odabultum	145	80 (43.2%)		
	Tuloworeda	132	49 (26.5%)		
	Gemachis	51	27 (14.6%)		
Total	Mieso	56	29 (15.7%)		
		384	185 (48.5%)		

In the current investigation, 185 (48.2%) of the total inspected organs were rejected for a variety of reasons. The liver was the organ that was condemned the most (27.3%), followed by the lung (14.3%), heart (4.68%), heart, and kidney (1.8%). The most frequent macroscopic lesions on the liver were fascioliasis (14.6%), cirrhosis (6.25%), hydatid cyst (3.13%), calcification

(2.6%), and abscess (0.78%). With rates of 6.5%, 4.69%, 2.34%, and 0.78%, respectively, pneumonia, emphysema, abscess, and hydatid cyst all contributed to the rejection of the lung. 1.5% of hearts and 2.6% of hearts, respectively, were condemned due to hydatid cyst and pericarditis. 1.8% of renal cases resulted in death from hydro nephrosis (Table 3).

Table 3. Distribution of condemned organ with respect to lesion found.

Organ Condemned	No. of Inspected	Lesion found	Condemnation (%)	Proportion (%)
Liver	105	Fasciolosis	56 (14.6)	53.3
		Hydatid cyst	12 (3.13)	11.4
		Cirrhosis	24 (6.25)	22.9
		Abscess	3 (0.78)	2.9
		Calcification	10 (2.6)	9.5
		Total	105 (27.3)	
Lung	55	Hydatid cyst	25 (6.25)	45.5
		Pneumonia	18 (4.09)	32.7
		Abscess	3 (0.78)	5.5
		Emphysema	9 (2.34)	16.4
		Total	55 (14.3)	
Heart	18	Pericarditis	10 (2.6)	55.6
		Cysticercusbovis	2 (0.5)	11.1
		Hydatid cyst	6 (1.5)	33.3
		Total	18 (4.68)	
Kidney	7	Hydronephrosis	7 (1.8)	100
		Total	7 (1.8)	
Grand total	185	Total	185 (48.2)	100

Liver condemnation was more common in cattle with medium body condition (4.69%), good body condition (8.07%), and poor body condition (4.69%), in that order. In a similarly, medium-conditioned cattle had greater rates of lung, heart, and kidney rejections than good- and poor-conditioned

animals. Adult cattle have greater liver, lung, heart, and kidney rejection rates than young cattle, and male cattle shows greater rejection rates than female calves for all examined organs (Table 4).

Table 4. Distribution of condemned organ with regard to age, sex, body condition score.

Variables	No. of Examined	Liver	Lung	Condemned organ Heart	Kidney	Total
Age	Young	151	40	15	4	60 (39.7%)
	Adult	233	65	40	14	125 (53.6)
BCS	Poor	32	18	7	3	28 (87%)
	Medium	207	56	32	10	112 (49.3%)
	Good	145	31	16	5	55 (37.9%)
Sex	Female	51	15	7	4	27 (52.9%)
	Male	333	90	48	14	158 (47.4%)

In the current study, the estimated annual direct financial loss as a result of organ condemnation was 138784.00 ETB (Table 3). Liver condemnation accounts for the highest

percentage of economic losses, or 27.3% of the overall losses of \$11998.00 ETB, followed by the lungs, heart, and kidney, which account for all of the direct economic losses (Table 5).

Table 5. The rejection rate and average price of organ condemned in the study area.

Condemned Organ	Rate of condemned Organ (%)	Average price in ETB	Total loss in ETB
Liver	27.3%	75	111998.00
Lung	14.3%	20	15644.00
Heart	4.68%	30	7696.00
Kidney	1.8%	35	3446.00
Total price (ETB)			138784.00

4. Discussion

The present study examined the factors for organ condemnation and the financial harm they generated in 384 cattle assessed at the Chiro municipal abattoir during the study period. According to this study, the most common reasons for organ condemnation in cattle slaughtered at the Chiro municipal abattoir were fasciolosis, hydatid cyst, pneumonia, *Cysticercus bovis*, cirrhosis, hydronephrosis, calcification, emphysema, pericarditis, and abscess. Out of 185 tested organs, the liver, the lung, the heart, and the kidney were all found to be defective in 27.3%, 14.3%, 4.68%, and 1.8% of cases, respectively. This result is consistent with earlier research from the Jimma Municipal slaughterhouse and Nekemte Municipal abattoir, which was published by Nebyou et al., and Yellow et al., [33, 44].

In the current study, 27.3% of the liver was deemed defective due to different abnormalities discovered during the postmortem examination. Fasciolosis was found to be one of the major reasons for liver rejection in this study with a prevalence of 14.6%, which is lower than the prevalences of 21.1 and 20.18% reported by Shitaye et al., [41], from the Hawasa Municipal abattoir and Yellow et al., [44] from the Dessie Municipal abattoirs, respectively. The findings of the current study outperform those of Hassan, [23]'s study from Iran (7.9%) and Alembrian and Hylagebriel, [3] study from the municipal slaughterhouse in Adigrat (9.26%). The discrepancy in climatic and ecological variables, such as height, rainfall, and temperature, between this research and the aforementioned findings may be primarily blamed for the difference in the rejection rate of liver owing to fasciolosis.

Findings of 6.25 percent liver condemnation owing to cirrhosis are close to Raji et al., [39] s 8.4 percent findings at Zara slaughterhouse. However, this figure is greater than the 1.1% reported by Yifat, [45] in Gonder, while the present study's liver condemnation rate owing to cirrhosis was lower than the 16.5% reported by Nurit et al., [36] in Kombolcha. Compared to the report of 1.9% in Tanzania by Nongoad, calcification was shown to be the cause of 2.6% of liver condemnation from the entire studied organ (2010). The rejection rate of the liver owing to hydatidosis in the current study is 3.13%, which is close to studies done by Dawit, [14] from the municipal slaughterhouse in Adigrat. Bzuayehu, [10] Reported migration rates from Harar of 3.6% and 3. 62%,

respectively. However much lower than the reports of Zelalem et al., [46] 31.7% from Addis Abeba and Mihret et al., [32], 33.33% from Dire Dawa. In contrast to Ahmedullah et al., [1], who reported a 3.8% liver condemnation rate in Bangladesh, the current study found a 0.78% liver condemnation owing to abscess.

14.3% of the lungs that were examined were condemned for different reasons. Hydatidosis and pneumonia were the main contributors to lung rejection among these causes. The lung condemnation rate (6.25%) in the current research is lower than the 19.37% reported by Gebremeskel and Kalayou, [19] from Mekelle. However, this rate is better than the 1.42% reported by Regassa et al., [40] for Ghana. This variance in hydatidosis prevalence may result from various animal husbandry practices, backyard killing of animals, improper disposal of diseased carcasses, the existence of stray dogs and their interactions with animals, as well as other factors by Mesale et al., [30]. Pneumonia-related lung death in the current research was 4.09%. It was higher than the amount (0.14%) that Clement et al. reported in cattle slain at the Zango slaughterhouse [13]. Some many factors, such as stress factors like exposure to environmental dust or exhaustion from long journeys made by pastoral livestock in search of pasture and water or when animals are taken to livestock markets or abattoirs, as well as parasitism, may explain the different prevalence of pneumonic lungs. [9].

Based on the most recent study, the heart's overall condemnation rate is 4.69%. Pericarditis, hydatid cysts, and *Cysticercus bovis* accounted for 2.6%, 1.5%, and 0.5% of all heart rejections in this research, respectively. This compares to studies by Genet, [20] from the Gonder slaughterhouse with 7.8% and 9.4%, respectively, which is incredibly lower. However, it was greater than the findings of Lati et al., [27] and the Nekemte Municipal Abattoir's 0.18% prevalence, respectively. *Cysticercus bovis* account for two cases (0.5%) of heart condemnation, which is somewhat equivalent to the prevalence of 0.23% reported by Basem, from Egypt. This acquired value, however, is less than that published by Garedaghi, [18], who indicated an Iranian prevalence of 3%. This difference in the reported prevalence rate may be caused by a number of variables, including variations in meat consumption habits, patient knowledge of the disease's clinical picture, personnel and environmental hygiene, control measures, and eradication campaigns. Additionally, according to the current study, fewer kidneys were condemned owing to

hydronephrosis than was reported by Fufa and Debele, [17] in their report from the municipal slaughterhouse at Walaita Sodo. The variance in disease prevalence and animal care practices at various research sites may be the reason of the variability in the rejection rate of organs with connected to various causes.

In this study, the annual direct financial loss attributable to the condemnation of organs at the Chiro Municipal abattoir was estimated to be 138,784.00 Ethiopian Birr, which is higher than the direct financial loss analysis estimated by Bekele et al. [8] from the Hossana Municipal abattoir and Alemayehu et al. [22] from the Luna export abattoir, which estimated a total financial loss of about 88,806. The current finding, however, is significantly lower than the estimate made by Haimanot et al., [22], who estimated yearly losses of 109,492,725.5 ETB as a result of organ condemnation from calves at the Dire Dawa Municipal abattoir. The difference in the financial loss estimated in various abattoir and/or parts of Ethiopia would be due to the variations in the prevalence of disease, mean annual number of cattle slaughtered in the different abattoirs and also the variation in the retail market price of organs [6].

5. Conclusions and Recommendations

The purpose of the current study, which was carried out in the municipal abattoir in Chiro, was to determine the main causes that organs are rejected and to estimate the direct financial loss connected with condemned organs. Based on the findings of the current study, the condemnations of organs at the Chiro Municipal Abattoir in Eastern Ethiopia were linked to a number of disease issues. 185 (48.2%) of the 384 animals that were investigated overall had postmortem abnormalities of various kinds. Organ condemnation rates were greater for the liver and lower for the lungs, hearts, and kidneys, correspondingly. Fasciolosis was the major pathological disease that led to the entire condemnation of the organs, followed by cirrhosis, calcification, emphysema, pericarditis, and hydronephrosis. Age group, provenance, and animal body condition all significantly affect organ condemnation rates ($P < 0.05$). According to estimates, edible organ condemnation results in annual financial losses of 138,784.00 ETB.

Based on the above conclusion the following recommendations are forwarded:

- 1) It is important to study the epidemiology of the disorders that lead to organ condemnation and implement control measures.
- 2) It is important to encourage the building of abattoirs with the requisite disposal pits and to dispose of rejected organs promptly and correctly.
- 3) The management of condemned carcasses and organs should be taught to meat inspectors and slaughterhouse staff.
- 4) In cooperation with the government, the veterinary extension service should be strengthened to raise community understanding of the disease's prevention and control.

References

- [1] Ahmedullah, F. M., M. G. Akbor, M. M. Kaider and K. Hossain, (2007): Pathological investigation of liver of the slaughtered buffaloes in Barisal district, Bangladesh. *J. Vet. Med.* 5: 81-85.
- [2] Alemayehu, R., Nebyou M, Bekele M, Desta B, DessieSh, Etana D, Fufa A, Eystein S (2012). Major causes of organs and carcass condemnation in small ruminants slaughtered at Luna Export Abattoir, Oromia Regional State, Ethiopia. *Prev. Vet. Med.* 110 (2): 139-148.
- [3] Alembrhan, A., and Hylagebriel, T (2013): Major cause of organ condemnation in cattle slaughtered at Adigrat municipal abattoir, northern Ethiopia, *Veterinary World* 6 (10): 734-738.
- [4] Alton, G. D., Lpeah, D., Bateman, K. J., McNab, W. B and Berk, O. (2010): Factors associated with whole condemnation rates in provincially inspected abattoir in Ontario 2001-2007: Implication for food animal syndromic surveillance. *BMC Vet. Res.*, 6 (42): 312-345.
- [5] Amene, F., Eskindir, L., Dawit, T. (2012): Cause Rate and Economic Implication of Organ Condemnation of Cattle Slaughtered at Jimma Municipal Abattoir, Southwestern Ethiopia. *Glob. Vet.* 9: 396-400.
- [6] Arbabi, M., Hooshy H (2006). Survey of Regions might have accounted for variation of the Echinococcosis and Hydatidosis in Kashan Region, Prevalence in different areas of a country Central Iran. *Iran. J. Public Health.* 35: 75-81.
- [7] Basem, R., Abdo, N., Amal, A., Sayed, M Asmaa, A. Hussein, A Moshen A and Arafa, I (2009): Occurrence of Cysticercosis in cattle and buffaloes and Taeniasaginata in man in Assiut Governance of Egypt. *Animal Research Institute, Assiut, Egypt. Dept. of Animal Hygiene and Zoonoses. Fac. Vet. Med. Assiut University, Egypt.*
- [8] Bekele, Ch., Sissay M, Mulugeta D (2014). On Farm Study of Bovine Fasciolosis in Lemo District and its economic loss due to liver condemnation at Hossana Municipal abattoir, Southern Ethiopia. *Int. J. Curr. Microbiol. App. Sci.* 3 (4): 1122-1132.
- [9] Benard, L. M., E. N. Hezron and Esron, D. K. (2011): Slaughter stock abattoir survey of carcasses and organ/offal condemnations in Arusha region, northern Tanzania. *Tropical Animal Health and Production*, 43: 857-64.
- [10] Bzuayehu, L., Tsegalem, A., Befikadu, U., Ayalew, N., Abebe, A. (2014): Prevalence of Bovine Hydatidosis and its Economic significance in Harar Municipality Abattoir, Eastern Ethiopia. *Ameri. Euro J scient res.* 9: 143-149.1.
- [11] CDLFO, (2015): Chiro district Livestock and Fishery Office.
- [12] Central Statistical Authority (CSA), (2013): Agricultural statistical report, Addis Ababa, Ethiopia.
- [13] Clement, B. J., I. J. Alawa and B. L. Joseph, 2010. A 6- years survey of pathological conditions of slaughtered animals at Zango abattoir in Zaria, Kudu state, Nigeria. *Tropical Animal Health and Production.* 43: 127-131.
- [14] Dawit A., Tewodros, A. and Tefera, K. (2017): The Prevalence of Bovine Hydatidosis among Slaughtered Cattle at DebreBerhan Municipal Abattoir, North Shewa Zone, Ethiopia. *JVetSciMed.* 5 (1): 5.

- [15] Elsa, L., Sofia, V., Elvira, S., Maria, M., Mendes, G., and André, M. (2012): Factors Influencing Livestock Productivity. Environmental Stress and Amelioration in Livestock Production, DOI: 10.1007/978-3-642-29205-7-2.
- [16] Food and Agriculture Organization (FAO) (1994): Manual of meat inspection for developing countries.
- [17] Fufa, A., and Debele, H. (2013): Major Causes of Organ Condemnation for Cattle and its Financial Impact at Wolaita Sodo Municipality Abattoir, Southern Ethiopia. 11 (6): 730-734.
- [18] Garedaghi, Y., Saber, R. A. P., and Khosroshahi, S. M. (2012): Prevalence of Bovine Cysticercosis of Slaughtered Cattle in Meshkinshahr Abattoir, Iran. Journal of Animal and Veterinary Advances. 11: 785-788.
- [19] Gebremeskel, B. and Kalayou, S. (2009): Prevalence, viability and fertility study of bovine cystic echinococcosis in Mekelle city, Northern Ethiopia. Revue méd. Vét. 160 (2): 92- 97.
- [20] Genet, M., Guadu, T., Basaznew, C., and Mersha, B. (2012): Pathological Conditions Causing Organ and Carcass Condemnation and Their Financial Losses in Cattle Slaughtered in Gondar, Northwest Ethiopia. 4 (6): 200-208.
- [21] Gracey, JF., Collins, OS., and Huey, RJ. (1999): Meat hygiene, 10th ed. London, Bailliere Tindall. 223-260.
- [22] Haimanot, D., Munera, A., Hawi, J., Tilahun, Z and Girma, K, (2015) Major cause of organ condemnation and its economic loss in cattle slaughtered at Dire Dawa Municipal abattoir, eastern, Ethiopia. Academic journal of animal disease. (4) 3: 118-123.
- [23] Hassan, B., Mohammad, A. and Mehrab, K. (2012) A retrospective study of abattoir condemnation due to parasitic infections: economic importance in Ahwaz, southwestern Iran. J. Parasitol. 98 (5): 954—957.
- [24] Jatenie, J., Mahendra, P., Rahman, T. (2014): Investigation into major causes of organs condemnation in bovine slaughtered at Adama municipal abattoir and their economic importance. Haryana. Vet. 53: 139-143.
- [25] KAOAR, (2016): Kombolcha agricultural office of animal resource recording sheet found.
- [26] Kidanu, L. (2011): Major parasitic case of organ condemnation in cattle and its economic importance at Jimma Municipal Abattoir, DVM Thesis, School of Veterinary Medicine, Jimma University College of Agriculture and Veterinary Medicine, Jimma Ethiopia.
- [27] Lati E, Biresaw S, Berhanu S, Eyob H (2015). Causes of organ condemnation, its public health and financial significance in Nekemte municipal abattoir, Wollega, Western Ethiopia. J. Vet. Med. Anim. Health. 7: 205-214.
- [28] Manyazewal, A., Zeleke, MG., Tsegaye, T. (2014): Economic Significance of Fasciolosis at Mettu Municipal Abattoir, Southwest Ethiopia. J. Adv. Vet. Res. 4 (2): 53-59.
- [29] Mellau, LS., Longa, HE., and Karimuribu, ED. (2010): A Slaughter house survey of liver lesion in slaughter cattle, sheep and goats at Arusha, Tanzania. Vet. J. 3: 179-188.
- [30] Mesele, A., Solomon T, Desie S (2013). Cystic Echinococcosis of Cattle in Jimma Municipal Abattoir, South West Ethiopia. Glob. Vet. 11 (6): 771-775.
- [31] Mezegebu, Y. (2003): Major cause of organ condemnation in ruminants slaughtered at Gonder Abattoir, North Western Ethiopia. DVM thesis, Faculty of Veterinary Medicine, Addis Ababa University, Debre-Zeit, Ethiopia.
- [32] Miheret, M., M. Biruk, T. Habtamu and K. Ashwani, 2013. Bovine hydatidosis in Eastern Part of Ethiopia. MEJS. 5 (1): 107-114.
- [33] Nebyou, M., Debela, A., Solomon, K and Tesema, T. (2014): Major Causes of Organs and Carcass Condemnation in Cattle Slaughtered at Nekemte Municipality Abattoir, East Wollega, Ethiopia. Glob Vet. 13: 278-284.
- [34] Nicholson, MJ., and Butter, MH. (1986): A guideline to condition scoring of Zebu cattle. International Livestock Center for Africa, Addis Ababa, Ethiopia.
- [35] Nongaond, E. D., Mellau, L. S. and (2010): Slaughter house survey liver lesions in slaughtered cattle, sheep and goats at Arasha, Tanzania. Journal of Vet Sci. 3: 179-188.
- [36] Nurit, M., Zerihun, H. and Serkalem, M. (2012): Major Cause of Liver Condemnation and Associated Financial Loss at Kombolcha Elmore Abattoir, South Wollo, Ethiopia. Europ. J. Appl. Sci. 4 (4): 140-145.
- [37] Ogunrinade, A., and Ogunrinade, BI. (1980): Economic importance of bovine fasciolosis in Nigeria. Trop. Anim. Health Prod. 12: 155-160.
- [38] Pace, J. E., and Wakeman, D. L. (2003): Determining the age of cattle by their teeth animal science department institute of food and agricultural sciences (IFAS). 312-348.
- [39] Raji, M. A., Solam, S. O., and Ameh, JM. (2010): Pathological condition and lesions observed in slaughtered cattle at Zaria abattoir. J. Clin. Pathol. Forensic Med. 1: 9-12.
- [40] Regassa, A., Abbu, F., mulugeta, A., and Megersa, B. (2009): Major Metacestodes in cattle at Wolaita Sodo municipal abattoir, Southern Ethiopia: Prevalence, cyst viability, organ distribution and socio-economic implications. Trop An. Health and Production. 41: 1495-1502.
- [41] Shitaye, M., Tilaye, S., and Fanos, T. (2016): Major causes of organ condemnation and associated financial loss in cattle slaughtered at Hawassa Municipal Abattoir, Ethiopia. J. Vet. Med. Anim. Health. 8 (10): 150-156.
- [42] Thrusfield, M. (2007): Veterinary Epidemiology. (3rd edition.). Blackwell Science.
- [43] Wale, T., Tirfayehu, M., Assefa, A. (2017): Major cause of organ condemnation and financial significance in cattle slaughtered at Jimma municipal Abattoir, southwestern, Ethiopia. Int. J. Adv. Res. Bio. Sci. (4) 3: 32-33.
- [44] Yellow, Zewdu, M., and Wedajo, M. (2016): Major causes of organ condemnation and associated financial loss in cattle slaughtered at Hawassa Municipal Abattoir, Ethiopia. J. Vet. Med. Anim. Health. 8 (7): 56-63.
- [45] Yifat, D., Gedefaw, D., and Desie, S. (2011): Major cause of organ condemnation and financial significance of cattle slaughtered at Gondar ELLFORA abattoir. Global Veterinarian. 7 (5): 487-490.
- [46] Zelalem, F., Tadele, T., Zelalem, N., Chanda, M and Nigatu, Kebede, G. (2012): Prevalence and characterization of hydatidosis in animals slaughtered at Addis Ababa abattoir, Ethiopia. J. Parasitol. Vector. Biol. 4 (1): 1-6.